

LB 01.1 | Subcutaneous Prophylaxis with the Anti-TFPI Monoclonal Antibody Concizumab in Hemophilia A and Hemophilia A/B with Inhibitors: Phase 2 Trial Results

J. Astermark¹; P. Angchaisuksiri²; G. Benson³; G. Castaman⁴; P. Chowdary⁵; H. Eichler⁶; V. Jiménez-Yuste⁷; K. Kavakli⁸; T. Matsushita⁹; L. Hvitfeldt Poulsen¹⁰; J. Oldenburg¹¹; A.D. Shapiro¹²; A. Wheeler¹³; S. Zupancic-Salek¹⁴; G. Young¹⁵

¹Lund University, Center for Thrombosis and Hemostasis, Malmö, Sweden, ²Ramathibodi Hospital, Mahidol University, Department of Medicine, Division of Hematology, Bangkok, Thailand, ³Belfast Health and Social Care Trust, Department of Hematology, Belfast, United Kingdom, ⁴Careggi University Hospital, Center for Bleeding Disorders and Coagulation, Department of Oncology, Florence, Italy, ⁵Royal Free London NHS Foundation Trust, Katharine Dormandy Hemophilia Centre and Thrombosis Unit, London, United Kingdom, ⁶Saarland University and University Hospital, Institute of Clinical Hemostaseology and Transfusion Medicine, Homburg, Germany, ⁷La Paz University Hospital, Haematology Department, Madrid, Spain, ⁸Ege University Children's Hospital, Department of Hematology, Izmir, Turkey, ⁹Nagoya University Hospital, Department of Transfusion Medicine, Nagoya, Japan, ¹⁰The Hemophilia Centre, Department of Haematology, Aarhus, Denmark, ¹¹University Hospital Bonn, Department of Immunohematology and Department of Molecular Hemostasis, Bonn, Germany, ¹²Indiana Hemophilia & Thrombosis Center, Inc., Indianapolis, United States, ¹³Vanderbilt University Medical Center, Department of Pathology, Microbiology and Immunology, Nashville, United States, ¹⁴University Hospital Centre Zagreb, Medical School of Osijek and Medical School of Zagreb, Department of Haematology, Haemophilia and Thrombosis Unit, Zagreb, Croatia, ¹⁵University of Southern California, Keck School of Medicine, Los Angeles, United States

Background: We present results from the concizumab explorer4 (NCT03196284; randomized) and explorer5 (NCT03196297; single-arm) phase 2 trials (main phase: 24 weeks).

Aims: To assess the safety, efficacy and immunogenicity of once-daily, subcutaneous concizumab prophylaxis in patients with hemophilia A/B with inhibitors (HAWI/HBWI) and hemophilia A without inhibitors (HA), and to establish a dose regimen for the phase 3 trials.


Methods: Informed consent and ethics committee approval were obtained. Thirty-six HA, nine HAWI and eight HBWI patients were exposed to concizumab. Patients received 0.15 mg/kg concizumab with potential dose escalation to 0.20 and 0.25 mg/kg. The number of adverse events (AEs) was evaluated, as well as coagulation-related parameters (D-dimer and prothrombin fragment 1+2 [F1+2]). Immunogenicity was assessed by the number of anti-drug antibodies (ADAs). Concizumab and free tissue factor pathway inhibitor (TFPI) plasma levels were measured by ELISA, and peak thrombin generation (TG) potential by a standardized assay.

Results: Clinical proof-of-concept (CPoC) for once-daily subcutaneous concizumab prophylaxis was established, there were no

thromboembolic events, no AE-related withdrawals, no safety concerns with breakthrough-bleed treatment, and annualized bleeding rates after 24 weeks were estimated at 3.0–7.0, with all patients choosing to continue to the extension phase in both trials. Concizumab exposure, free TFPI reduction and TG potential were similar across all hemophilia patients (Fig.1A–C), and concizumab exposure was associated with normalized TG potential (Fig.1D). With increasing concizumab concentration, elevated F1+2 were detected (Fig.1E), and there was a tendency for elevated D-dimers (Fig.1F). Three patients in each trial had ADA-positive tests, with no apparent impact on any measured parameters.

Conclusions: Phase 2 trial results confirmed CPoC, with no unexpected safety signals and have guided selection of the phase 3 dosing regimen. Further development of concizumab as a prophylactic treatment for all hemophilia patients is supported, including HBWI patients who currently have no prophylactic options.

Figure 1 Mean plots for patients with last dose level of concizumab 0.15 mg/kg (A–C) and for all dose levels (D–F) in the main phase of explorer4 (HAWI/HBWI) and explorer5 (HA without inhibitors) of (A) concizumab plasma concentration vs time; (B) free TFPI vs time; (C) peak thrombin generation potential vs time; (D) peak thrombin generation potential vs concizumab plasma concentration; (E) prothrombin fragments 1+2 vs concizumab plasma concentration; and (F) D-dimers vs concizumab plasma concentration, by hemophilia type.



Baseline free TFPI, mean (standard deviation): 96.3 (11.1) ng/mL. Normal range of peak thrombin generation (lower; upper): 26; 147 nmol/L. HA, hemophilia A; HAWI, hemophilia A with inhibitors; HBWI, hemophilia B with inhibitors; TFPI, tissue factor pathway inhibitor

FIGURE 1

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2019 The Authors. *Research and Practice in Thrombosis and Haemostasis* published by Wiley Periodicals, Inc on behalf of International Society on Thrombosis and Haemostasis.

LB 01.2 | First-in-human Evidence of Durable Therapeutic Efficacy and Safety of AAV Gene Therapy Over Three-years with Valoctocogene Roxaparvovec for Severe Haemophilia A (BMN 270-201 Study)

K.J. Pasi¹; S. Rangarajan²; N. Mitchell³; W. Lester⁴; M. Laffan⁵; B. Madan⁶; E. Symington⁷; B. Kim³; X. Yang³; J. Quinn³; J. Henshaw³; C. Vettermann³; G.F. Pierce⁸; W.Y. Wong³

¹Royal London Haemophilia Centre Barts Health NHS Trust, London, United Kingdom, ²University Hospital Southampton, Southampton, United Kingdom, ³BioMarin Pharmaceutical Inc., Novato, United States, ⁴University Hospitals Birmingham NHS Foundation Trust, Birmingham, United Kingdom, ⁵Centre for Haematology, Imperial College London, London, United Kingdom, ⁶Guys & St Thomas' NHS Foundation Trust, London, United Kingdom, ⁷Cambridge University Hospitals NHS Foundation Trust, Cambridge, United Kingdom, ⁸Consultant, La Jolla, United States

Background: Gene therapy is increasingly viewed as a viable treatment option for hemophilia A (HA), using AAV mediated Factor VIII (FVIII) gene transfer. Interim Phase 1/2 data from valoctocogene roxaparvovec (AAV5-hFVIII-SQ) have shown promising results. Outstanding questions for all AAV gene therapies relate to clinical effectiveness and durability.

Aims: Assessment of long-term safety, efficacy, and durability of AAV5-hFVIII-SQ in a Phase 1/2 clinical study.

Methods: Adult male study participants with severe HA received a single intravenous dose of AAV5-hFVIII-SQ at 6×10^{13} vg/kg (n=7) or 4×10^{13} vg/kg (n=6).

Results: All study participants demonstrated clinically meaningful FVIII activity levels with reductions in bleeds and FVIII usage (Figure 1). Following withdrawal from prophylaxis, annualised bleeding rate (ABR) declined from pre-treatment mean by 96% at year three in 6×10^{13} vg/kg participants, and 92% at year two in 4×10^{13} vg/kg participants (Figure 2). FVIII levels reported by chromogenic assay correspond with the continued absence of target joints and target joint bleeds from years two through three. Expression levels over time are determined to decline as a function of both time post-administration and level of




FIGURE 1 Factor VIII levels by dose cohort over time




FIGURE 2 Annualised bleeding rates and FVIII usage by dose cohort over time

expression achieved, nearing a plateau of expression in year three. Accordant modelling projections conservatively estimate the persistence of bleeding control for at least eight years post-administration and longer if expression plateaus are maintained, as observed in prior AAV gene therapy studies (Figure 1). The safety profile of valoctocogene roxaparvovec remains favourable and unchanged, with no inhibitor development or ALT elevations beyond year one. Detailed durability, safety, and efficacy data will be shared at ISTH.

Conclusions: Gene transfer with valoctocogene roxaparvovec has resulted in substantial and sustained FVIII activity levels, clinically relevant reductions in self-reported bleeding episodes, and significant reductions in FVIII replacement infusions for up to three years post-dosing.

*This study was approved by a recognized medical ethics committee at each participating site. All participants provided informed consent.

LB 01.3 | A Multicenter Trial of Vena Cava Filters in Severely-Injured Patients

K.-M. Ho^{1,2,3}; R. Sudhakar⁴; S. Honeybul⁵; R. Zellweger⁴; B. Wibrow⁶; J. Lipman^{7,8}; A. Holley^{7,8}; A. Kop⁹; E. Geelhoed¹⁰; T. Corcoran¹¹; P. Misur¹²; C. Edibam¹³; R. Baker¹⁴; J. Chamberlain²; C. Forsdyke⁴; F. Rogers¹⁵

¹University of Western Australia, Intensive Care Medicine, Perth, Australia, ²Royal Perth Hospital, Intensive Care Medicine, Perth, Australia, ³Murdoch University, School of Veterinary & Life Sciences, Perth, Australia, ⁴Royal Perth Hospital, Trauma, Perth, Australia, ⁵Royal Perth Hospital, Neurosurgery, Perth, Australia, ⁶Sir Charles Gairdner Hospital, Intensive Care Medicine, Perth, Australia, ⁷Royal Brisbane and Women's Hospital, Intensive Care Medicine, Brisbane, Australia, ⁸University of Queensland, Intensive Care Medicine, Brisbane, Australia, ⁹Centre for Implant Technology and Retrieval Analysis, Medical Engineering & Physics, Perth, Australia, ¹⁰University of Western Australia, School of Allied Health, Perth, Australia, ¹¹University of Western Australia, School of Medicine and Pharmacology, Perth, Australia, ¹²Royal Perth Hospital, Radiology, Perth, Australia, ¹³Fiona Stanley Hospital, Intensive Care Medicine, Perth, Australia, ¹⁴Murdoch University, Centre for Thrombosis and Haemostasis, Perth, Australia, ¹⁵University of Pennsylvania College of Medicine, Surgery, Lancaster, United States

Background: It is not known whether early placement of an inferior-vena-cava filter reduces death or pulmonary embolism in severely-injured patients with a contraindication to anticoagulant prophylaxis.

Aims: We aimed to assess whether insertion of a retrievable vena cava filter within 72 hours of trauma admission reduces mortality or symptomatic pulmonary embolism.

Methods: In this multicenter randomized-controlled-trial, 240 severely-injured patients - with an Injury-Severity-Score (ranges between 0 and 75 with higher scores indicating more severe injury) >15 and contraindications to anticoagulants - were randomly allocated to receive a vena cava filter or no filter within the first 72 hours of admission. The primary outcome was a composite of 90-day mortality or symptomatic pulmonary embolism; secondary outcomes included symptomatic pulmonary embolism between day 8 and 90 for the subgroup of patients who survived at least 7 days and did not receive prophylactic anticoagulation within 7 days after injury. All patients received screening lower limb ultrasound at 2 weeks, and mandatory computed-tomography pulmonary-angiography when prespecified criteria were met.

Results: The median age and Injury-Severity-Score of the patients were 39 years and 27, respectively. Compared to no filter, early filter placement did not significantly reduce the primary outcome (13.9% vs 14.4%, respectively; hazard ratio 0.99, 95% CI 0.51-1.94, P=0.98). In those who did not receive anticoagulant prophylaxis within 7 days after injury, none in the filter group compared to five in the control group developed pulmonary embolism, including one fatal event (0% vs 14.7%, respectively, relative risk: 0, 95%CI 0.00-0.55). Entrapped thrombus within the filter was found in six patients (4.9%).

Conclusions: Early prophylactic placement of a vena cava filter after major trauma did not reduce the primary composite outcome of 90-day mortality or symptomatic pulmonary embolism.

(ACTRN trial number: 12614000963628)

LB 01.4 | Magnetic Resonance Direct Thrombus Imaging Can Safely Rule Out Recurrent Ipsilateral Deep Vein Thrombosis of the Leg - The Theia study

L.F. van Dam¹; C.E.A. Dronkers²; G. Gautam³; Å. Eckerbom³; W. Ghanima⁴; J. Gleditsch⁵; A. von Heijne³; H.M.A. Hofstee²; M.M.C. Hovens⁶; S. Kolman⁷; L.J.M. Kroft⁸; A.T.A. Mairuhu⁹; M. Nijkeuter¹⁰; M.A. van de Ree⁷; C.J. van Rooden¹¹; R.E. Westerbeek¹²; J. Westerink¹⁰; E. Westerlund³; M.V. Huisman¹; F.A. Klok¹

¹Leiden University Medical Center, Thrombosis and Hemostasis, Leiden, the Netherlands, ²Haaglanden Medical Center, Internal Medicine, The Hague, the Netherlands, ³Karolinska Institutet, Danderyd Hospital, Clinical Sciences, Stockholm, Sweden, ⁴Ostfold Hospital Trust, Internal Medicine, Ostfold, Norway, ⁵Ostfold Hospital Trust, Radiology, Ostfold, Norway, ⁶Rijnstate Hospital, Vascular Medicine, Arnhem, the Netherlands, ⁷Diakonessenhuis, Vascular Medicine, Utrecht, the Netherlands, ⁸Leiden University Medical Center, Radiology, Leiden, the Netherlands, ⁹Haga Teaching Hospital, Internal Medicine, The Hague, the Netherlands, ¹⁰University Medical Center Utrecht, Vascular Medicine, Utrecht, the Netherlands, ¹¹Haga Teaching Hospital, Radiology, The Hague, the Netherlands, ¹²Deventer Hospital, Radiology, Deventer, the Netherlands

Background: The diagnosis of recurrent ipsilateral deep vein thrombosis (DVT) of the leg is challenging because of frequent persistent




FIGURE 1 Coronal MRDTI images

intravascular abnormalities after previous DVT. Magnetic Resonance Direct Thrombus Imaging (MRDTI), a technique without intravenous contrast and with a 10-minute acquisition time, has been shown to reliably distinguish acute recurrent DVT from chronic DVT (Figure 1).

Aims: To evaluate the safety of ruling out recurrent ipsilateral DVT of the leg by normal MRDTI.

Methods: The Theia study is a multicentre, international diagnostic management study (NCT02262052) in patients with suspected acute recurrent ipsilateral DVT of the leg. Main exclusion criteria were concurrent suspected pulmonary embolism (PE) or previous ipsilateral DVT ≤6 months before presentation. The final treatment decision was based on the MRDTI result alone, which was performed within 24 hours of study inclusion. The primary endpoint was the 3-month incidence of venous thromboembolism (VTE) in patients with normal MRDTI at baseline. A VTE incidence with an upper limit of the 95% confidence interval (95%CI) below 6.6% was deemed safe. We also determined the interobserver agreement of MRDTI reading. All endpoints were adjudicated by an independent committee. The study was approved by all local IRBs. All patients provided written informed consent.

Results: Of 444 patients screened between May 2014 and March 2019, 305 (69%) were included (Figure 2). The baseline prevalence of recurrent DVT was 37%; thrombophlebitis was diagnosed in an additional 4.6%. The primary endpoint occurred in 2 of 121 (1.7% (95%CI 0.20-5.8)) patients with normal MRDTI who did not receive any form of (prophylactic) anticoagulation during follow-up; neither of these recurrences was fatal. The agreement between initial local reading and post-hoc central reading of the MRDTI images was excellent (kappa statistic 0.92).

Conclusions: Recurrent ipsilateral DVT can safely be ruled out by MRDTI alone, which can replace the often non-conclusive compression ultrasound in these patients.




FIGURE 2 Study flowchart


LB 01.5 | Rivaroxaban for the Treatment of Acute Venous Thromboembolism in Children

C. Male¹; A. Lensing²; D. Kubitzka²; A. Chan³; G. Kenet⁴; P. Massicotte⁵; G. Young⁶; A.C. Molinari⁷; U. Nowak-Göttl⁸; M. Crowther⁹; M. Prins¹⁰; P. Monagle¹¹; for the EINSTEIN-Jr. Phase 3 Investigators

¹Medical University of Vienna, Paediatrics, Wien, Austria, ²Bayer AG, Wuppertal, Germany, ³McMaster University, Children's Hospital, Hamilton, Ontario, Canada, ⁴Tel Aviv University, Sheba Medical Center, Institute of Thrombosis and Hemostasis, Tel Hashomer, Israel, ⁵University of Alberta, Department of Pediatrics, Edmonton, Canada, ⁶University of Southern California, Children's Hospital of Los Angeles, Los Angeles, United States, ⁷University of Genova, IRCCS Giannina Gaslini, Genova, Italy, ⁸University Hospital Schleswig-Holstein, Institute for Clinical Chemistry, Kiel, Germany, ⁹McMaster University, Department of Medicine, Hamilton, Ontario, Canada, ¹⁰Maastricht University Medical Center, Maastricht, the Netherlands, ¹¹University of Melbourne, Royal Children's Hospital, Melbourne, Australia

Background: Treatment of venous thromboembolism (VTE) in children is currently mostly based on adult data. Bodyweight-adjusted rivaroxaban regimens had previously been established for children (birth-17 years) to match the exposure range of young adults treated with rivaroxaban 20 mg once-daily.

Aims: To compare the efficacy and safety of rivaroxaban with standard anticoagulants (comparator) for treatment of acute VTE in children.



Population PK results for adult VTE patients treated with rivaroxaban 20 mg once-daily are shown as box whisker plot (5, 25, 50, 75, and 95 percentiles, extremes as open circles).

FIGURE 1 Individual population PK parameters for rivaroxaban at steady state as a function of bodyweight

Methods: 500 children (birth-17 years) with acute VTE (lower extremity 33.0%, cerebral veins 23.4%, lungs 16.0%, upper extremity 11.6%, jugular vein 10.4%) were randomized (2:1 ratio) to receive open-label rivaroxaban (tablets or suspension) in bodyweight-adjusted 20-mg equivalent dose regimens or comparator. The main treatment period was 3 months (1 month in children < 2 years with catheter-related VTE). Symptomatic recurrent VTE and bleeding were centrally assessed unaware of treatment assignment. End of treatment repeat imaging was obtained. Pharmacokinetic (PK) and pharmacodynamic (PD) data were collected.

Results: Recurrent VTE occurred in 4/335 (1.2%) rivaroxaban-recipients and in 5/165 (3.0%) comparator-recipients (hazard ratio 0.40, 95% confidence interval, 0.11 to 1.41). Repeat imaging showed an improved effect of rivaroxaban on thrombotic burden as compared with comparator (table). Clinically relevant bleeding occurred in 10 children (3.0%; all nonmajor) with rivaroxaban and in 3 children (1.9%; two major; 1 nonmajor) with comparator. Absolute/relative efficacy and safety estimates of rivaroxaban and comparator were similar to those obtained in adults. PK parameters for rivaroxaban confirmed plasma levels in children equal to those achieved in adult

TABLE 1 Change in thrombotic burden on repeat imaging at the end of the main treatment period as compared to the index event

Thrombotic burden on repeat imaging	Rivaroxaban (n=335)	Comparator (n=165)
Normalised	129 (38.5%)	43 (26.1%)
Improved	130 (38.8%)	75 (45.5%)
Uncertain	55 (16.4%)	28 (17.0%)
No relevant change	16 (4.8%)	13 (7.9%)
Deterioration	1 (0.3%)	1 (0.6%)
Symptomatic recurrent VTE	4 (1.2%)	5 (3.0%)

van Elteren test for comparison of ordered categories, $p=0.009$.

VTE trials that showed efficacy of rivaroxaban (figure). PK and PD were closely correlated. Clinical and PK/PD results were similar for rivaroxaban tablets and suspension.

Conclusions: In children with VTE, treatment with bodyweight-adjusted rivaroxaban (tablets or suspension) resulted in low

recurrence risk and reduced thrombotic burden without increased bleeding compared to standard anticoagulants. Our results are comparable to those obtained in adults with VTE treated with rivaroxaban.

- A**
Angchaisuksiri P. LB 01.2
Astermark J. LB 01.2
- B**
Baker R. LB 01.5
Benson G. LB 01.2
- C**
Castaman G. LB 01.2
Chamberlain J. LB 01.5
Chan A. LB 01.1
Chowdary P. LB 01.2
Corcoran T. LB 01.5
Crowther M. LB 01.1
- D**
Dronkers C.E.A. LB 01.3
- E**
Eckerbom Å. LB 01.3
Edibam C. LB 01.5
Eichler H. LB 01.2
- F**
Forsdyke C. LB 01.5
- G**
Gautam G. LB 01.3
Geelhoed E. LB 01.5
Ghanima W. LB 01.3
Gleditsch J. LB 01.3
- H**
Henshaw J. LB 01.4
Ho K.-M. LB 01.5
Hofstee H.M.A. LB 01.3
Holley A. LB 01.5
Honeybul S. LB 01.5
Hovens M.M.C. LB 01.3
- Huisman M.V. LB 01.3
Hvitfeldt Poulsen L. LB 01.2
- J**
Jiménez-Yuste V. LB 01.2
- K**
Kavakli K. LB 01.2
Kenet G. LB 01.1
Kim B. LB 01.4
Klok F.A. LB 01.3
Kolman S. LB 01.3
Kop A. LB 01.5
Kroft L.J.M. LB 01.3
Kubitza D. LB 01.1
- L**
Laffan M. LB 01.4
Lensing A. LB 01.1
Lester W. LB 01.4
Lipman J. LB 01.5
- M**
Madan B. LB 01.4
Mairuhu A.T.A. LB 01.3
Male C. LB 01.1
Massicotte P. LB 01.1
Matsushita T. LB 01.2
Misur P. LB 01.5
Mitchell N. LB 01.4
Molinari A.C. LB 01.1
Monagle P. LB 01.1
- N**
Nijkeuter M. LB 01.3
Nowak-Göttl U. LB 01.1
- O**
Oldenburg J. LB 01.2
- P**
Pasi K.J. LB 01.4
- Pierce G.F. LB 01.4
Prins M. LB 01.1
- Q**
Quinn J. LB 01.4
- R**
Rangarajan S. LB 01.4
Rogers F. LB 01.5
- S**
Shapiro A.D. LB 01.2
Sudhakar R. LB 01.5
Symington E. LB 01.4
- V**
Van Dam L.F. LB 01.3
Van de Ree M.A. LB 01.3
Van Rooden C.J. LB 01.3
Vettermann C. LB 01.4
Von Heijne A. LB 01.3
- W**
Westerbeek R.E. LB 01.3
Westerink J. LB 01.3
Westerlund E. LB 01.3
Wheeler A. LB 01.2
Wibrow B. LB 01.5
Wong W.Y. LB 01.4
- Y**
Yang X. LB 01.4
Young G. LB 01.1, LB 01.2
- Z**
Zellweger R. LB 01.5
Zupancic-Salek S. LB 01.2